

# Product Catalogue

**Advancing  
Science**

**For A Healthier  
Future**



For further information please contact us at :



**BERNOFARM**  
PHARMACEUTICAL COMPANY

Ruko Puri Mansion Blok C15, Jln. Lingkar Luar Barat  
RT 2 RW 1. Kembangan Selatan, Jakarta barat  
Phone : 021-29025795

Websites : [www.bernofarm.com](http://www.bernofarm.com)



Certificate No : ID12/02206



Certificate No : ID12/02214



Certificate No : ID12/02214

Being a pharmaceutical company **RANKED 8th among all multinational and local companies in Indonesia (IQVIA Q2 2024)** is an encouragement for Bernofarm to continue developing its capacity and capabilities. Starting from a small drugstore ; then developed to a pharmaceutical company that has its own manufacture – not only for Bernofarm products but also for other companies (toll in). Bernofarm covers the market for ethical products (branded & generic), OTC (over the counter) products and/or supplements, as well as international business. Now Bernofarm has more than 500+ registered products distributed in Indonesia and several countries.

1971 : Bernofarm was established in Surabaya, East Java – Indonesia

1976 : Expanded from 20 employees, gradually over than 3000 people at present. Our manufacturing site was moved to a strategic location at East Java.

1991 : Obtained Good Manufacturing Practice (GMP) from Ministry of Health (MOH) Indonesia

2009 : ISO 9001, continuously updated ISO 14001, continuously updated

2012 : OHSAS 18001, continuously updated

2019 : Halal Assurance System, continuously updated

2020 : ISO 45001, continuously updated

**NOW** : Accredited GMP from Malaysia, Mongolia, Maldives, Myanmar, Cambodia, Fiji, Republic Dominican, Timor Leste, etc.



For further information please contact us at :



**BERNOFARM**  
PHARMACEUTICAL COMPANY

Ruko Puri Mension Blok C15, Jln. Lingkar Luar Barat  
RD2 RW 1 Kembangan Selatan, Jakarta barat  
Phone : 021-29025795

Websites : [www.bernofarm.com](http://www.bernofarm.com)



Certificate No : ID12/02206



Certificate No : ID12/02205



Certificate No : ID12/02214

# Content

## Introduction

### A. Gastrointestinal Drugs

KLIRAN	01
PANTOTIS	02
ESOPRAZ	03
INHIPRAZ	04
SULCOLON	05

### B. Blood and Cardiovascular Agents

LOVASK	07
CANDEPRESS	08
TRANXA	09
EPODION 2000 IU	10

### E. Anti-Infectives

ECOTRIXON	11
SULBACEF	12
TOCEF	13
MACEF	14
MEROPEX	15
BERNOFLOX	16
LEVOVID	17
MOXIVID	18
ZISTIC	19
TRACHON	20
FUNGISTOP	21

### F. Anesthetics

TIOPOL	23
LIDODEX	24
BUCAIN SPINAL HEAVY	25
BELNIUM	26

### G. Dermatological Agent

BERSOL	28
BENOSON-G	29
BENOSON-M	30

### H. Analgesic and Anti-Inflammation

PARAMOL	32
TROLAC	33
ANAFEN	34
METHYLON	35
SULCOLON	36

### I. Nerves and Radiocontrast

BRALIN	38
BRALIN	39
BERTHYCO	40
HEXIOL	41

### J. Over The Counter Medicine (Kids)

BIOLYSIN	43
BIOLYSIN KIDS	44
BIOLYSIN KIDS C 100	45
BIOLYSIN EMULSION	46
BIOLYSIN PLATINUM	47
BIOLYSIN SMART	48
BONE KIDZ	49

### K. Over The Counter Medicine (Adult)

VITALONG C	51
VITALONG C	52
VITALONG C + ZINC	53
VITALONG C + ZINC	54
GREAT D	55
FITERGY	56
BIOLYSIN	57

# Gastrointestinal Drugs







## **Composition**

Ondansetron HCl Dihydrate ~ Ondansetron 8 mg

## **Dosage Form**

Film coated caplet

## **Indication**

Nausea and Vomiting after chemotherapy, radiotherapy, or surgery

## **Dosage and Administration**

Adult:

8 mg twice daily up to 5 days

Children:

4 mg twice daily up to 5 days

## **Storage Condition**

Store below 30° C, protect from light

## **Packaging**

Box, 3 strips @ 10's

## **MOQ (Box)**

1.667

## **Shelf Life**

5 years (Subjects to change)

# PANTOTIS

ATC level : A02BC



## Composition

Pantoprazole Sodium Sesquihydrate (Lyophilized) ~Pantoprazole  
40 mg / vial

## Dosage Form

Powder for injection

## Indication

Increased gastric acid production, gastro-esophagus reflux disease,  
Zollinger-Ellison syndrome

## Dosage and Administration

Adult:

Gastric and duodenal ulcer, reflux oesophagitis: 40 mg once daily  
up to 7 days

Zollinger-Ellison syndrome: 80 mg daily up to 7 days

## Storage Condition

Store below 25 C and dry place

After reconstitution

Solutions stable for 12 hours at temperature below 25° C

## Packaging

Box, 1 vial @ 40 mg

## MOQ (Box)

4.273

## Shelf Life

3 years (Subjects to change)

# ESOPRAZ

ATC level : A02BC



## Composition

Esomeprazole Sodium Sterile Lyophilized ~ Esomeprazole 40 mg / vial

## Dosage Form

Powder for injection

## Indication

Increased gastric acid production, gastro-esophagus reflux disease, Zollinger-Ellison syndrome

## Dosage and Administration

GERD and gastric ulcer 20-40 mg once a day  
for rebleeding after endoscopy ; 80 mg for 30 minute and then  
8 mg / hour for 3 days

## Storage Condition

Store below 30° C protect from light  
After reconstitution Solution stable for 12 hours at temperature  
below 30° C

## Packaging

Box, 1 vial @ 40 mg

## MOQ (Box)

8.474

## Shelf Life

2 years (Subjects to change)



ATC level : A02BC



### **Composition**

Lansoprazole Pellet ~ Lansoprazole 30 mg

### **Dosage Form**

Delayed release capsule

### **Indication**

Increased gastric acid production, gastro-esophagus reflux disease, Zollinger-Ellison syndrome.

### **Dosage and Administration**

Adult:

30 mg once daily 4 up to 8 weeks

### **Storage Condition**

Store below 25° C and dry place, protect from light

### **Packaging**

Box, 1 strip @ 10's

### **MOQ (Box)**

10.000

### **Shelf Life**

2 years (Subjects to change)



### **Composition**

Sulfasalazine 500 mg

### **Dosage Form**

Enteric coated caplet

### **Indication**

Ulcerative colitis, Crohn's Disease, Rheumatoid Arthritis

### **Dosage and Administration**

Adult:

2-4 caplets daily in divided doses

Children:

40 - 60 mg/kg body weight daily, maintenance 20 - 30 mg/kg body weight

### **Storage Condition**

Store below 30° C, protect from light

### **Packaging**

Box, 10 strips @ 10's

### **MOQ (Box)**


3.000

### **Shelf Life**

2 years (Subjects to change)



# Blood and Cardiovascular Agents





## Composition

Amlodipine Besilate ~ Amlodipine 10 mg

## Dosage Form

Tablet

## Indication

Hypertension, coronary artery disease, Prinzmetal angina

## Dosage and Administration

Adult:

Initial dose 5 mg once daily, maintenance dose 5 mg to 10 mg once daily

## Storage Condition

Store below 30° C

## Packaging

Box, 3 blister @ 10's

## MOQ (Box)

3.333

## Shelf Life

5 years (Subjects to change)



# CANDEPRESS

ATC level : C09CA



## Composition

Candesartan Cilexetil 8 mg

## Dosage Form

Tablet

## Indication

Hypertension, heart failure, coronary heart disease

## Dosage and Administration

Hypertension:

adult 4 mg a day, max 16 mg a day

elderly >75 year; 2 mg once a day

Heart failure:

initial dose 4 mg a day, Up-titration to the target dose of 32 mg once daily or the highest tolerated dose

## Storage Condition

Store below 30° C, protect from light

## Packaging

Box, 3 blisters @ 10's

## MOQ (Box)

3.333

## Shelf Life

2 years (Subjects to change)



ATC level : B02AA



### **Composition**

Tranexamic Acid 50 mg / ml

### **Indication**

Local fibrinolysis, abnormal bleeding during surgery, hereditary angioneurotic edema, menorrhagia, bleeding after dental surgery

### **Dosage Form**

Injection

### **Dosage and Administration**

5-10 ml administered via im/iv after operation or divided into 2 doses

10 - 50 ml administered via infusion

### **Storage Condition**

Store below 30° C, protect from light

### **Packaging**

Box, 5 ampoules @ 5 ml


### **MOQ (Box)**

3.774

### **Shelf Life**

5 years (Subjects to change)

# Anti-Infectives





# ECOTRIXON

ATC level : J01DD



## Composition

Ceftriaxone Sodium Sesquaterhydrate~ Ceftriaxone 1 g / vial

## Dosage Form

Powder for injection

## Indication

Respiratory tract infection, soft and skin tissue infection, sexually transmitted disease, blood infection, nerve infection, prophylaxis antibiotic

## Dosage and Administration

Adult (>12 yr):

1-2 g once daily, administered every 24 hours. For severe infections maximum daily dose can be increased up to 4 g per day

Neonates (<14 days):

20 - 50 mg/kg body weight per day

Infants and Children (15 days - 12 yr):

20 - 80 mg/kg body weight per day

## Storage Condition

Store below 25 C and dry place

After reconstitution

Solutions stable for 6 hours at temperature below 30° C or for 24 hours at 5° C

## Packaging

Box, 1 vial @ 1 g + solvent @ 10 ml

## MOQ (Box)

8.403

## Shelf Life

5 years (Subjects to change)



# SULBACEF

ATC level : J01DD



## Composition

Cefoperazone Sodium and Sulbactam Sodium ~ Cefoperazone 500 mg dan Sulbactam 500 mg

## Dosage Form

Powder for injection

## Indication

Respiratory tract infection, urinary tract infection, intra-abdominal infection, skin and soft tissue infection

## Dosage and Administration

Adult:

mild/moderate: 2-4 g every 12 hours

severe: up to 8 g every 12 hours

## Storage Condition

Store below 30° C, protect from light

After reconstitution

Solution stable for 7 days at temperature 2 - 8° C and for 24 hours at below 25° C

## Packaging

Box, 1 vial @ 1 g + aqua pro Injection @ 5 ml

## MOQ (Box)

4.363

## Shelf Life

3 years (Subjects to change)



### Composition

Cefixime Trihydrate ~ Cefixime Anhydrous 200 mg

### Dosage Form

Film coated caplet

### Indication

Urinary tract infection, respiratory tract infection, typhoid fever, sexually transmitted disease

### Dosage and Administration

Adult & Children > 30 kg:

mild/moderate: 50 - 100 mg twice daily

severe: up to 200 mg twice daily

### Storage Condition

Store below 25° C and dry place

### Packaging

Box, 1 blister @ 10's

### MOQ (Box)

4.000

### Shelf Life

3 years (Subjects to change)



### Composition

Cefixime Trihydrate ~ Cefixime 100 mg / 5 ml

### Dosage Form

Dry syrup

### Indication

Urinary tract infection, respiratory tract infection, typhoid fever, sexually transmitted disease

### Dosage and Administration

Children :

1,5 - 3 mg / kg can be increased to 6 mg/kg

typhoid: 10 -15 mg/kg once a day for 2 weeks

### Storage Condition

Store below 25° C and dry place

After reconstitution can be stored for up to 7 days at room temperature or under refrigeration

### Packaging

Box, 1 bottle @ 30 ml

### MOQ (Box)

10.000

### Shelf Life

3 years (Subjects to change)



## Composition

Cefepime HCl Arginine ~ Cefepime 1 g / vial

## Dosage Form

Powder for injection

## Indication

Respiratory tract infection, urinary tract infection, intra-abdominal infection, gynaecological infection, and skin infection

## Dosage and Administration

UTI (mild to moderate): 0,5 -1 g 2 times daily for 7 up to 10 days

UTI (severe): 2 g, 2 times daily for 10 days

Pneumonia (moderate - severe): 1 g - 2 g, 2 times daily for 10 days

Skin and skin structure infection: 2 g every 12 hours for 10 days

Empiric treatment in febrile neutropenia: 2 g every 8 hours for 7 days

Intra-abdominal infection with complication: 2 g, 2 times daily for 7 up to 10 days

Children: The above dosage can be used for children that have weight over 40 kg,

For children under 40 kg: 50 mg/kg every 12 hours. (beside febrile neutropenia)

Febrile neutropenia 40 kg: 50 mg/kg every 8 hours.

## Storage Condition

Store below 25° C and dry place, protect from light

After reconstitution

Solution stable for 24 hours at temperature below 25° C, or for 7 days in a refrigerator (2 - 8°) C

## Packaging

Box, 1 vial @ 1 g + solvent @ 10 ml

## MOQ (Box)

2.724

## Shelf Life

3 years (Subjects to change)



# MEROPEX

ATC level : J01DH



## Composition

Meropenem Trihydrate Bufferred Sodium Carbonate ~ Meropenem  
1 g / vial

## Dosage Form

Powder for injection

## Indication

Respiratory tract infection, urinary tract infection, intra-abdominal infection joint infection, gastrointestinal gynaecological infection, skin infection, blood infection, nosocomial infection

## Dosage and Administration

Adult:

Pneumonia, UTI, gynaecological infections: 500 mg IV every 8 hours  
Pneumonia, peritonitis, presumed infections in neutropenic patients and septicemia: 1 g every 8 hours

Meningitis: 2 g every 8 hours

Children:

> 3 months up to 12 yr: 10 - 20 mg/kg every 8 hours  
> 50 kg body weight: 40 mg/kg every 8 hours

## Storage Condition

Store below 30 °C, protect from light

After reconstitution

Solutions stable for 8 hours at temperature below 25° C or for 48 hours at 2 - 8° C

## Packaging

Box, 1 vial @ 1 g

## MOQ (Box)

3.500

## Shelf Life

2 years (Subject to change)



# BERNOFLOX

ATC level : J01MA



## Composition

Ciprofloxacin HCl Monohydrate ~ Ciprofloxacin 500 mg

## Dosage Form

Film coated tablet

## Indication

urinary tract infection, respiratory tract infection, skin and soft tissue infection, bone and joint infection, gastrointestinal tract infection

## Dosage and Administration

Adult:

Mild/moderate: 2 times 250 - 500 mg daily

Severe/Chronic: 2 times 500 - 750 mg daily

## Storage Condition

Store below 30° C, protect from light

## Packaging

Box, 2 strips @ 10's

## MOQ (Box)

5.000

## Shelf Life

5 years (Subjects to change)



## Composition

Levofloxacin Hemihydrate ~ Levofloxacin 500 mg

## Dosage Form

Film coated caplet

## Indication

Respiratory tract infection, skin infection, urinary tract infection

## Dosage and Administration

Adult:

250 - 750 mg once daily for 3 - 14 days

## Storage Condition

Store below 30° C

## Packaging

Box, 1 blister @ 10 's

## MOQ (Box)

10.000

## Shelf Life

5 years (Subjects to change)



### Composition

Moxifloxacin HCl ~ Moxifloxacin 400 mg

### Dosage Form

Film coated tablet

### Indication

Respiratory tract infection, Pneumonia, Tuberculosis

### Dosage and Administration

Acute exacerbation of chronic bronchitis: 400 mg once daily for 5 up to 10 days

CAP: 400 mg once daily for 10 days

Acute sinusitis: 400 mg once daily 10 days

### Storage Condition

Store below 30° C

### Packaging


Box, 1 strips @ 10's

### MOQ (Box)

1.000

### Shelf Life

5 years (Subjects to change)



## Composition

Azithromycin Dihydrate ~ Azithromycin 500 mg

## Dosage Form

Film coated caplet

## Indication

Respiratory tract infection, skin infection, sexually transmitted disease

## Dosage and Administration

Adult:

500 mg daily for 3 days

Sexually transmitted disease the dose is 1000 mg as single oral dose

Children > 6 months:

10 mg/kg dose on day 1, 5 mg/kg day 2 to 5

## Storage Condition

Store below 30° C

## Packaging

Box, 1 strip @ 10's

## MOQ (Box)

8.500

## Shelf Life

4 years (Subjects to change)



### **Composition**

Itraconazole Pellet ~ Itraconazole 100 mg

### **Dosage Form**

Capsule

### **Indication**

Fungal infection except meningitis

### **Dosage and Administration**

Adult:

100 - 200 mg once to twice daily

### **Storage Condition**

Store below 30° C, protect from light

### **Packaging**

Box, 1 strip @ 10's

### **MOQ (Box)**

10.000

### **Shelf Life**

3 years (Subjects to change)





### **Composition**

Griseofulvin 500 mg

### **Dosage Form**

Film coated tablet

### **Indication**

Hair and scalp fungal infection

### **Dosage and Administration**

Adult:

500 - 1000 mg once or divided dose daily

Children:

10 mg/kg body weight once or divided dose daily

### **Storage Condition**

Store below 30° C

### **Packaging**

Box, 10 strips @ 10's


### **MOQ (Box)**

1.000

### **Shelf Life**

4 years (Subjects to change)

# Anesthetics





## Composition

Thiopental Sodium With Sodium Carbonate ~ Thiopental Sodium  
0,5 g / vial

## Dosage Form

Powder for injection

## Indication

General anesthesia, drug-resistant epilepsy

## Dosage and Administration

Adult:

50 to 75 mg at intervals of 20 to 40 seconds

## Storage Condition

Store below 25 C and dry place

After reconstitution

Can be stored for up to 24 hours at temperatures below 25° C

## Packaging

Box, 5 vial @ 0,5 g

## MOQ (Box)

742

## Shelf Life

5 years (Subjects to change)



### **Composition**

Lidocaine HCl Monohydrate ~ Lidocaine HCl 50 mg / ml  
and Dextrose 5%

### **Dosage Form**

Injection

### **Indication**

Spinal anesthesia and epidural anesthesia

### **Dosage and Administration**

1 - 2 ampoule (maximum dose is 2 ampoules)

### **Storage Condition**

Store below 30°C, protect from light

### **Packaging**

Box, 5 ampoules @ 2 ml

### **MOQ (Box)**

9.302

### **Shelf Life**

4 years (Subjects to change)

# BUCAIN SPINAL HEAVY

ATC level : N01BB



## Composition

Bupivacaine HCl Monohydrate ~ Bupivacaine HCl 5 mg; Dextrose Anhydrate 80 mg / ml

## Dosage Form

Injection

## Indication

Spinal anesthesia and epidural anesthesia

## Dosage and Administration

Lower-limb, urological and perineal surgery:  
7.5 - 15 mg, 2 - 3 hours

Lower abdominal surgery  
15 - 20 mg, 1.5 - 2 hours

## Storage Condition

Store below 30°C, protect from light

## Packaging

Box, 5 ampoules @ 4 ml

## MOQ (Box)

2.325

## Shelf Life

5 years (Subjects to change)

# BELNIUM

ATC level : M03AC



## Composition

Rocuronium Bromide 10 mg / ml

## Dosage Form

Injection

## Indication

Facilitate rapid intubation sequence and tracheal intubation, and to provide muscle paralysis

## Dosage and Administration

Adult:

IV bolus inj/continuous infusion Tracheal intubation 0.6 mg/kg.

Rapid sequence induction of anesthetic 1 mg/kg or 0.6 mg/kg

Maintenance: 0.15 mg/kg. Long-term inhalation anesthetic 0.075-0.1 mg/kg

## Storage Condition

Store in temperature between 2 - 8°C, protect from light and do not freeze this product

## Packaging

Box, 1 vial @ 5 ml


## MOQ (Box)

4.716

## Shelf Life

5 years (Subjects to change)

# Dermatological Agent





### **Composition**

Clobetasol Propionate 0,5 mg per g

### **Dosage Form**

Cream

### **Indication**

Skin inflammation (Ultra-High potency)

### **Dosage and Administration**

Adult, elder, children over 1 year:

Apply thinly and gently rub in using only enough to cover the entire affected area once or twice a day until improvement occurs

### **Storage Condition**

Store below 30°C, protect from light

### **Packaging**

Box, 1 tube @ 15 g

### **MOQ (Box)**

7.894

### **Shelf Life**

5 years (Subjects to change)



# BENOSON-G

ATC level : D07CC



## Composition

Betamethasone Valerate ~ Betamethasone 1 mg;  
Gentamicin Sulfate ~ Gentamicin 1 mg / g

## Dosage Form

Cream

## Indication

Bacterial skin infection accompanied by inflammation

## Dosage and Administration

Apply a thin layer to the afflicted area 2 times a day, in the morning and night

\*note: frequency of can be adjusted according to doctor's instruction

## Storage Condition

Store below 25°C, protect from light

## Packaging

Box, 1 tube @ 10 g

## MOQ (Box)

11.764

## Shelf Life

5 years (Subjects to change)



### Composition

Betamethasone Valerate ~ Betamethasone 1 mg;  
Miconazole Nitrate 20 mg / g

### Dosage Form

Cream

### Indication

Fungal skin infection accompanied by inflammation

### Dosage and Administration

Apply a thin layer to the afflicted area 2 times a day

### Storage Condition

Store below 30°C, protect from light

### Packaging

Box, 1 tube @ 5 g


### MOQ (Box)

23.076

### Shelf Life

3 years (Subjects to change)

# Analgesic and Anti-Inflammation



# PARAMOL

ATC level : N02BE



## Composition

Paracetamol 1 g / 100 ml

## Dosage Form

Infusion

## Indication

Fever, Mild Pain

## Dosage and Administration

Adolescents and Adult (> 50kg):

1g up to 4 times a day

Children (33 kg - 50 kg):

15 mg/kg per administration up to 4 times a day

## Storage Condition

Store below 30°C

## Packaging

Box, 1 bottle @ 100 ml

## MOQ (Box)

29.126

## Shelf Life

2 years (Subjects to change)



## Composition

Ketorolac Trometamol 30 mg / ml

## Dosage Form

Injection

## Indication

Moderate to severe pain

## Dosage and Administration

initial dose 10 mg followed by 10 - 30 mg every 4 - 6 hours

max dose 90 mg / day

Elderly, patient w/ renal function impairment, patient <50 kg Max: 40 mg. Max duration of treatment: 2 days.

## Storage Condition

Store below 30°C

## Packaging

Box, 5 ampoules @ 1 ml

## MOQ (Box)

4.348

## Shelf Life

4 years (Subjects to change)

# ANAFEN

ATC level : M01AE



## Composition

Ibuprofen 100 mg / 5 ml

## Dosage Form

Suspension

## Indication

Mild to moderate pain

## Dosage and Administration

Adult:

mild to moderate pain : 200 mg 3-4 times a day

Children:

fever and mild to moderate pain:

1 - 2 years: 50 mg 3 -4 times a day

3 - 7 years: 100 mg 3-4 times a day

8 - 12 years: 200 mg 3 -4 times a day

should not be given to children below one year

## Storage Condition

Store below 30°C

## Packaging

Box, 1 bottle @ 60 ml

## MOQ (Box)

9.677

## Shelf Life

3 years (Subjects to change)

# METHYLON

ATC level : H02AB



## Composition

Methylprednisolone 4 mg

## Dosage Form

Tablet

## Indication

Accute and chronic inflammation

## Dosage and Administration

Adult

Initially, 4-48 mg daily up to 160 mg daily should not be used in long term

Children

Use the lowest effective dose for the shortest possible duration

## Storage Condition

Store below 30°C, protect from light

## Packaging

Box, 10 strips @ 10's

## MOQ (Box)

5.000

## Shelf Life

2 years (Subjects to change)

# SULCOLON

ATC level : A07EC



## Composition

Sulfasalazine 500 mg

## Dosage Form

Enteric coated caplet

## Indication

Ulcerative colitis, Crohn's Disease, Rheumatoid Arthritis

## Dosage and Administration

Adult:

2-4 caplets daily in divided doses

Children:

40 - 60 mg/kg body weight daily, maintenance 20 - 30 mg/kg body weight

## Storage Condition

Store below 30°C, protect from light

## Packaging

Box, 10 strips @ 10's

## MOQ (Box)


3.000

## Shelf Life

2 years (Subjects to change)



# Nerves and Radiocontrast





### Composition

Citicoline 500 mg

### Dosage Form

Tablet

### Indication

Nerve repair after head trauma and open-angle glaucoma

### Dosage and Administration

500 mg, 1 tablet daily, with or without food.

### Storage Condition

Store below 30° C, protect from light

### Packaging

Box, 3 blisters @ 10's

### MOQ (Box)

3.333

### Shelf Life

3 years (Subjects to change)



### Composition

Citicoline Sodium ~ Citicoline 125 mg / ml

### Dosage Form

Injection

### Indication

Nerve repair after head trauma

### Dosage and Administration

For acute stage :

Usually 250 – 500 mg, 1 – 2 times daily by intravenous drip or intravenous injection.

For chronic stage :

Usually 100 – 300 mg, 1 – 2 times daily by intravenous or intramuscular injection.

Dosage may be increased according to condition.

### Storage Condition

Store below 30°C, protect from light

### Packaging

Box, 5 ampoules @ 8 ml

### MOQ (Box)

2.353

### Shelf Life

3 years (Subjects to change)



## Composition

Citicoline Sodium ~ Citicoline 125 mg / ml

## Dosage Form

Injection

## Indication

Nerve repair after head trauma

## Dosage and Administration

For acute stage :

Usually 250 – 500 mg, 1 – 2 times daily by intravenous drip or intravenous injection.

For chronic stage :

Usually 100 – 300 mg, 1 – 2 times daily by intravenous or intramuscular injection.

Dosage may be increased according to condition.

## Storage Condition

Store below 30°C, protect from light

## Packaging

Box, 5 ampoules @ 4 ml

## MOQ (Box)

4.651

## Shelf Life

5 years (Subjects to change)



### **Composition**

Mecobalamin 500 mcg

### **Dosage Form**

Capsule

### **Indication**

Peripheral neuropathies

### **Dosage and Administration**

1 capsule 3 times a day

### **Storage Condition**

Store below 30°C, protect from light

### **Packaging**

Box, 10 blisters @ 10's

### **MOQ (Box)**

3.000

### **Shelf Life**

2 years (Subjects to change)





## Composition

Iohexol ~ Iodine 350 mg I / mL

## Dosage Form

Injection

## Indication

Radio-contrast agent for radiographic procedures

## Dosage and Administration

The dosage depends on the type of investigation and the technique used. Usually the same iodine concentration and volume is used as for other iodinated X-ray contrast media in current use.

Adequate hydration should be assured before and after administration as for other contrast media.

## Storage Condition

Store below 30°C, protect from light

## Packaging

Box, 1 bottle @ 50 ml


## MOQ (Box)

1.960

## Shelf Life

2 years (Subjects to change)

# Over The Counter Medicine (OTC)



# BIOLYSIN

ATC level : A11A

## Composition

L-Lysine HCl 300 mg  
Vitamin A 1000 IU  
Vitamin B1 3 mg  
Vitamin B2 2 mg  
Vitamin B3 10 mg  
Vitamin B5 3 mg  
Vitamin B6 5 mg  
Vitamin B12 6 mcg  
Vitamin D3 200 IU  
Javanese turmeric extract 25 mg  
Honey 400 mg

## Dosage Form

Syrup

## Indication

Improve appetite and fulfill vitamin needs

## Dosage and Administration

1 tea spoon (5 mL), once daily

## Storage Condition

Store below 30°C, protect from light

## Packaging

Box, 1 bottle @ 60 mL

Box, 1 bottle @ 100 mL

## MOQ (Box)

19.672 (For bottle of 60 mL)

11.881 (For bottle of 100 mL)

## Shelf Life

2 years (Subjects to change)



# BIOLYSIN KIDS

ATC level : A11A



## Composition

Vitamin A 1500 IU  
Vitamin B1 1.4 mg  
Vitamin B2 1/6 mg  
Vitamin B6 1.6 mg  
Vitamin C 60 mg  
Vitamin D3 400 IU  
Vitamin E 5 mg  
Niacinamide 10 mg  
Calcium pantothenate 3 mg  
L-Lysine 50 mg

## Dosage Form

Chewable Tablet

## Indication

Help fulfill vitamin needs in children and during recovery.

## Dosage and Administration

1 chewable tablet, once daily

## Storage Condition

Store below 30°C, protect from light

## Packaging

Box, 1 bottle @ 30 chewable tablet

## MOQ (Box)

10.000

## Shelf Life

2 years (Subjects to change)



# BIOLYSIN KIDS C 100

ATC level : A11GA



## Composition

Vitamin C 100 mg

## Dosage Form

Chewable Tablet

## Indication

Help fulfill vitamin C needs and maintain immune system.

## Dosage and Administration

1 - 2 chewable tablet per day

## Storage Condition

Store below 30°C

## Packaging

Box, 1 bottle @ 30 chewable tablet

## MOQ (Box)

10.000

## Shelf Life

2 years (Subjects to change)

# BIOLYSIN EMULSION

ATC level : A11A



## Composition

Vitamin A 1000 IU  
Vitamin B1 0.6 mg  
Vitamin B2 0.6 mg  
Vitamin B6 0.6 mg  
Vitamin D3 200 IU  
Choline 75 mg  
DHA 10% 7 mg  
AA 40% 21 mg  
Cod Liver Oil 3 mg  
Mg Gluconate 50 mg  
Ca Lactate Gluconate 500 mg

## Dosage Form

Syrup

## Indication

Help maintain bone and teeth health. .

## Dosage and Administration

Children age 1 - 6 years old : 1 tablespoon (15 mL), once daily

Children age 6 - 12 years old : 1 tablespoon (15 mL), twice daily

## Storage Condition

Store below 30°C, protect from light

## Packaging

Box, 1 bottle @ 60 mL

Box, 1 bottle @ 250 mL

## MOQ (Box)

9.523 (For bottle of 60 mL)

4.705 (For bottle of 250 mL)

## Shelf Life

2 years (Subjects to change)

# BIOLYSIN PLATINUM

ATC level : A11A

## Composition

Vitamin A 1000 IU  
Vitamin B1 1.5 mg  
Vitamin B2 1.5 mg  
Vitamin B6 1.5 mg  
Vitamin B12 3 mcg  
Vitamin D3 200 IU  
Choline 75 mg  
L-Lysine HCl 100 mg  
DHA 10% 7 mg  
AA 40% 21 mg  
FOS 500 mg  
Curcuma xanthorrhiza 2 mg  
Fe 0.5 mg  
Folic acid 0.1 mg

## Dosage Form

Syrup

## Indication

Help fulfill vitamin needs and aid digestive system

## Dosage and Administration

Children age 1 - 6 years old : 1 tablespoon (15 mL), once daily

Children age 6 - 12 years old : 1 tablespoon (15 mL), twice daily

## Storage Condition

Store below 30°C, protect from light

## Packaging

Box, 1 bottle @ 250 mL

## MOQ (Box)

4.705

## Shelf Life

2 years (Subjects to change)



# BIOLYSIN SMART

ATC level : A11A



## Composition

AA Oil 21 mg  
Fish Oil (DHA 10%) 7 mg  
Choline 75 mg  
L-Lysine HCl 200 mg  
Vitamin A 1000 IU  
Vitamin D3 200 IU  
Vitamin B1 1.5 mg  
Vitamin B2 1.5 mg  
Vitamin B6 1.5 mg  
Vitamin B12 2 mcg

## Dosage Form

Syrup

## Indication

Help fulfill vitamin needs and improve brain function

## Dosage and Administration

Children age 1 - 6 years old : 1 tablespoon (15 mL), once daily

Children age 6 - 12 years old : 1 tablespoon (15 mL), twice daily

## Storage Condition

Store below 30°C, protect from light

## Packaging

Box, 1 bottle @ 60 mL

Box, 1 bottle @ 100 mL

## MOQ (Box)

19.354 (For bottle of 60 mL)

11.764 (For bottle of 100 mL)

## Shelf Life

2 years (Subjects to change)

# BONE KIDZ

ATC level :A11GC

## Composition

Lithothamnion sp. Extract  
(Calcium min. 45 mg) 150 mg  
Vitamin D3 25 IU  
FOS 50 mg

## Dosage Form

Chewable Tablet

## Indication

Help maintain bone and teeth health.

## Dosage and Administration

Children 2 - 4 years old : 1 - 2 chewable tablet per day

Children 5 - 12 years old : 2 - 4 chewable tablet per day

## Storage Condition

Store below 30°C, protect from light

## Packaging

Box, 1 bottle @ 30 chewable tablet

## MOQ (Box)


6.666

## Shelf Life

2 years (Subjects to change)



# Over The Counter Medicine (OTC)



# VITALONG C

ATC level :A11GA

## Composition

Vitamin C 500 mg

## Dosage Form

Time release capsule

## Indication

Increase stamina

## Dosage and Administration

1 capsule, once daily

## Storage Condition

Store below 30°C, protect from light

## Packaging

Box, 1 bottle @ 30 film coated caplet

## MOQ (Box)

6.666

## Shelf Life

2 years (Subjects to change)



# VITALONG C

ATC level :A11GA



## Composition

Vitamin C 500 mg

## Dosage Form

Time release capsule

## Indication

Increase stamina

## Dosage and Administration

1 capsule, once daily

## Storage Condition

Store below 30°C, protect from light

## Packaging

Box, 25 strips @ 4 film coated caplet

## MOQ (Box)

2.000

## Shelf Life

2 years (Subjects to change)



# VITALONG C+ZINC

ATC level :A11GB



## Composition

Vitamin C 500 mg

Zinc 5 mg

## Dosage Form

Time release capsule

## Indication

Maintain health and promote recovery

## Dosage and Administration

1 capsule, once daily

## Storage Condition

Store below 30°C, protect from light

## Packaging

Box, 1 bottle @ 30 film coated caplet

## MOQ (Box)

6.666

## Shelf Life

3 years (Subjects to change)

# VITALONG C + ZINC

ATC level :A11GB



## Composition

Vitamin C 500 mg

Zinc 5 mg

## Dosage Form

Time release capsule

## Indication

Maintain health and promote recovery

## Dosage and Administration

1 capsule, once daily

## Storage Condition

Store below 30°C, protect from light

## Packaging

Box, 25 strips @ 4 film coated caplet

## MOQ (Box)

2.000

## Shelf Life

3 years (Subjects to change)



### **Composition**

Vitamin D3 (Cholecalciferol) 1000 IU

### **Dosage Form**

Chewable Tablet

### **Indication**

Help fulfill Vitamin D needs, maintain bone health, and maintain immune system

### **Dosage and Administration**

Adult and children above 12 years old : 1 chewable tablet, once daily

### **Storage Condition**

Store below 30°C

### **Packaging**

Box, 1 bottle @ 100 chewable tablet

### **MOQ (Box)**

3.000

### **Shelf Life**

2 years (Subjects to change)

# FITERGY

ATC level : A11A

## Composition

L-Carnitine Fumarate 100 mg  
Panax Ginseng Extract 20 mg  
Ginger Extract 200 mg  
Caffeine 50 mg  
Vitamin B1 100 mg  
Vitamin B6 100 mg  
Vitamin B12 100 mcg  
Vitamin C 60 mg

## Dosage Form

Film coated caplet

## Indication

Increase stamina

## Dosage and Administration

1 film coated caplet, once daily

## Storage Condition

Store below 30°C, protect from light

## Packaging

Box, 25 strips @ 4 film coated caplet

Box, 1 bottle @ 30 film coated caplet

## MOQ (Box)

3.000 (For box of 25 strips)

10.000 (For box of 1 bottle)

## Shelf Life

2 years (Subjects to change)



# BIOLYSIN

ATC level : A11A



## Composition

Vitamin A Acetate 1500 IU  
Vitamin B1 1 mg  
Vitamin B2 0.8 mg  
Vitamin B6 0.8 mg  
Vitamin B12 3 mcg  
Vitamin C 50 mg  
Vitamin D2 400 IU  
Niacinamide 10 mg  
L-Lysine HCl 200 mg  
Ca Pantothenate 3 mg

## Dosage Form

Sugar coated tablet

## Indication

Improve appetite and fulfill vitamin needs

## Dosage and Administration

Adult : 1 sugar coated tablet, once daily

## Storage Condition

Store below 30°C

## Packaging

Box, 10 strips @ 10 sugar coated tablet

## MOQ (Box)

5.000

## Shelf Life

3 years (Subjects to change)

**For More Detailed Information, Scan Here**



Rukan Puri Mansion Blok C/15,  
West Jakarta, Jakarta, Indonesia



**+62898-6600-478 (Whatsapp / Telegram)**



**www.bernofarm.com**



*For further information please contact us at :*



**BERNOFARM**  
PHARMACEUTICAL COMPANY

Rukan Puri Mansion Blok C15, Jln. Lingkar Luar Barat  
RD 2 RW 1 Kembangan Selatan, Jakarta barat  
Phone : 021-29025795

Websites : [www.bernofarm.com](http://www.bernofarm.com)



Certificate No : ID12/02206



Certificate No : ID12/02205



Certificate No : ID12/02214